arXiv:1401.8150v1 [math.FA] 31 Jan 2014

ATOMIC DECOMPOSITIONS FOR OPERATORS IN REPRODUCING KERNEL HILBERT SPACES

LAURA GĂVRUŢA *

Abstract. In this paper, we give some results concerning atomic decompositions for operators on reproducing kernel Hilbert spaces, using frame theory techniques. We provide also generalizations for atomic decompositions of some theorems for reproducing kernel Hilbert spaces.
In particular, we obtain atomic decomposition results for operators on Bergman spaces and Fock spaces in a simple manner.

Keywords frames, atomic systems, Bergman spaces, reproducing kernel, Fock spaces

2010 Mathematics Subject Classification: 42C15, 30H20, 47B32

1 INTRODUCTION

In the following, we present some basic known facts about frames. Frames were introduced by R.J. Duffin and A.C. Schaffer [10] in 1952, in the context of nonharmonic Fourier series. In 1986, frames were brought to life by Daubechies, Grossman and Meyer in the fundamental paper [9].

We denote by $\mathcal{H}$ a separable Hilbert space and by $\mathcal{L}(\mathcal{H})$ the space of all linear bounded operators on $\mathcal{H}$.

DEFINITION 1 *A family of elements $\{f_n\}_{n=1}^{\infty} \subset \mathcal{H}$ is called a frame of $\mathcal{H}$ if there exists constants $A, B > 0$ such that*

$$A\|x\|^2 \leq \sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2 \leq B\|x\|^2, \quad x \in \mathcal{H}.$$

The constants A, B are called frame bounds.

If just the last inequality in the above definition holds, we say that $\{f_n\}_{n=1}^{\infty}$ is a *Bessel sequence*.

*The author was supported by the FWF project P 24986-N25

The operator

$$T : l^2 \to \mathcal{H}, \;\; T\{c_n\}_{n=1}^{\infty} := \sum_{n=1}^{\infty} c_n f_n$$

is called *synthesis operator(or pre-frame operator).* The adjoint operator is given by

$$\Theta = T^* : \mathcal{H} \to l^2, \;\; \Theta x = \{\langle x, f_n \rangle\}_{n=1}^{\infty}$$

and is called the *analysis operator.* By composing T with its adjoint T^* we obtain the *frame operator*

$$S : \mathcal{H} \to \mathcal{H}, \;\; Sx = TT^*x = \sum_{n=1}^{\infty} \langle x, f_n \rangle f_n.$$

The following Theorem is one of the most important result in frame theory because it gives the reconstruction formula.

THEOREM 1 *Let $\{f_n\}_{n=1}^{\infty} \subset \mathcal{H}$ be a frame for $\mathcal{H}$ with frame operator S. Then*

(i) S is invertible and self-adjoint;

(ii) every $x \in \mathcal{H}$ can be represented as

$$x = \sum_{n=1}^{\infty} \langle x, f_n \rangle S^{-1} f_n = \sum_{n=1}^{\infty} \langle x, S^{-1} f_n \rangle f_n. \tag{1.1}$$

For basic results on frame theory see the references [1], [3], [4], [8], [12], [13], [15].

The results in this paper are organized as follows. In Section 2, we recall the definition of an atomic system for operators and we complete a Theorem for the characterization of these systems, considered for the first time by the author in [7]. We rewrite this Theorem for a reproducing kernel Hilbert space. In Section 3 of this paper, we present some basic known facts about Bergman spaces, we give the atomic decomposition result for this case, for the standard weighted Bergman spaces and for Bergman spaces with Békollé-Bonami weights. In the last section of the paper, we give an atomic decomposition result for operators in Fock spaces.

2 ATOMIC DECOMPOSITIONS FOR OPERATORS

The first atomic decompositions results for holomorphic functions in Lebesgue spaces were obtained in 1980 by R.R. Coifman and R. Rochberg [5].

Let $\mathcal{H}$ be a separable Hilbert space and $\{f_n\}_{n=1}^{\infty} \subset \mathcal{H}$.

DEFINITION 2 *[7] Let $L \in \mathcal{L}(\mathcal{H})$. We say that $\{f_n\}_{n=1}^{\infty}$ is an atomic system for L if the following statements hold*

(i) the series $\sum_n c_n f_n$ converges for all $c = (c_n) \in l^2$;

(ii) there exists $C > 0$ such that for every $x \in \mathcal{H}$ there exists $a_x = (a_n) \in l^2$ such that $\|a_x\|_{l^2} \leq C\|x\|$ and $Lx = \sum_n a_n f_n$.

REMARK 1 *The condition (i) in Definition 2 actually says that $\{f_n\}_{n=1}^{\infty}$ is a Bessel sequence. (see Corollary 3.2.4 in [3])*

The above definition and the following main result (Theorem 3) presents a generalization of the concept of frames. The particular case of orthogonal projections was considered by H.G. Feichtinger and T. Werther [11].

The following Theorem for the existence of the atomic systems for an operator was proved in [7].

THEOREM 2 *Let $\mathcal{H}$ be a separable Hilbert space and $L \in \mathcal{L}(\mathcal{H})$. Then L has an atomic system.*

In the following Theorem, we present a characterization for atomic systems.

THEOREM 3 *Let $\{f_n\}_{n=1}^{\infty} \subset \mathcal{H}$. Then the following statements are equivalent*

(i) $\{f_n\}_{n=1}^{\infty}$ is an atomic system for L;

(ii) there exists $A, B > 0$ such that

$$A\|L^*x\|^2 \leq \sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2 \leq B\|x\|^2, \quad for\ any\ x \in \mathcal{H};$$

(iii) $\{f_n\}_{n=1}^{\infty}$ is a Bessel sequence and there exists a Bessel sequence $\{g_n\}_{n=1}^{\infty}$ such that

$$Lx = \sum_{n=1}^{\infty} \langle x, g_n\rangle f_n;$$

(iv) $\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2 < \infty$ and there exists a Bessel sequence $\{g_n\}_{n=1}^{\infty}$ such that

$$L^*x = \sum_{n=1}^{\infty} \langle x, f_n\rangle g_n, \forall x \in \mathcal{H}.$$

Proof. $(i) \iff (ii) \iff (iii)$ were proved in [7]

$(iv) \implies (ii)$ Using Cauchy-Schwartz inequality, we have:

$$\begin{aligned}\langle L^*x, L^*x\rangle &= \langle \sum_{n=1}^{\infty} \langle x, f_n\rangle g_n, L^*x\rangle \\ &= \sum_{n=1}^{\infty} \langle x, f_n\rangle \langle g_n, L^*x\rangle \\ &= |\sum_{n=1}^{\infty} \langle x, f_n\rangle \langle g_n, L^*x\rangle| \\ &\leq \left(\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2\right)^{1/2} \left(\sum_{n=1}^{\infty} |\langle g_n, L^*x\rangle|^2\right)^{1/2}.\end{aligned}$$

Since $\{g_n\}_{n=1}^{\infty}$ is a Bessel sequence, we obtain

$$\langle L^*x, L^*x\rangle \leq C\|L^*x\| \left(\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2\right)^{1/2}.$$

If $L^*x \neq 0$, we have $\|L^*x\|^2 \leq C\|L^*x\| \left(\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2\right)^{1/2}$ and from here we get

$$\frac{1}{C^2}\|L^*x\|^2 \leq \sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2$$

From hypothesis, $\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2 < \infty, \quad \forall x \in \mathcal{H}$ (actually $\{f_n\}_{n=1}^{\infty}$ is a Bessel sequence). From [3] we have that there exists $B > 0$ such that

$$\sum_{n=1}^{\infty} |\langle x, f_n\rangle|^2 \leq B\|x\|^2.$$

$(iii) \implies (iv)$

From (iii) we have $Lx = \sum_{n=1}^{\infty} \langle x, g_n\rangle f_n$.

For any $y \in \mathcal{H}$ we have

$$\begin{aligned}\langle L^*y, x\rangle = \langle y, Lx\rangle &= \langle y, \sum_{n=1}^{\infty}\langle x, g_n\rangle f_n\rangle \\ &= \sum_{n=1}^{\infty}\langle g_n, x\rangle\langle y, f_n\rangle \\ &= \sum_{n=1}^{\infty}\langle\langle y, f_n\rangle g_n, x\rangle \\ &= \langle\sum_{n=1}^{\infty}\langle y, f_n\rangle g_n, x\rangle\end{aligned}$$

which implies $L^*y = \sum_{n=1}^{\infty}\langle y, f_n\rangle g_n$ $\square$

DEFINITION 3 *We say that $\{f_n\}_{n=1}^{\infty}$ is a $L-$frame (or a frame for L) if there exists the constants $A, B > 0$ such that*

$$A\|L^*x\|^2 \leq \sum_{n=1}^{\infty}|\langle x, f_n\rangle|^2 \leq B\|x\|^2, \quad \forall x \in \mathcal{H}.$$

We consider now the reproducing kernel Hilbert space $(\mathcal{H}, K)$, with the kernel $K(z, \lambda) = K_\lambda(z)$, $K : \Omega \times \Omega \to \mathbb{C}$, where Ω is a given nonempty set.

We recall that a reproducing kernel Hilbert space $(\mathcal{H}, K)$ is a Hilbert space $\mathcal{H}$ of functions on Ω such that for every λ, K_λ belongs to $\mathcal{H}$ and for every $\lambda \in \Omega$ and every $f \in \mathcal{H}$, $f(\lambda) = \langle f, K_\lambda\rangle$.

Let $\Lambda = \{\lambda_n\}_{n=1}^{\infty} \subset \Omega$ be a subset of points with $\lambda_n \neq \lambda_m$, for $n \neq m$. In this case, with $f_n = \dfrac{K_{\lambda_n}}{\|K_{\lambda_n}\|}$, $n = 1, 2, \ldots$, we rewrite Theorem 3 as follows:

THEOREM 4 *The following statements are equivalent*

(i) $\left\{\dfrac{K_{\lambda_n}}{\|K_{\lambda_n}\|}\right\}_{n=1}^{\infty}$ *is an atomic system for L i.e.*

- *the series $\sum_{n=1}^{\infty} c_n \dfrac{K_{\lambda_n}}{\|K_{\lambda_n}\|}$ converges for all $\{c_n\} \in l^2$; and*
- *there exists $C > 0$ such that for every $f \in \mathcal{H}$ there exists $a_f = (a_n) \in l^2$ such that $\|a_f\|_{l^2} \leq C\|f\|$ and $Lf = \sum_{n=1}^{\infty} a_n \dfrac{K_{\lambda_n}}{\|K_{\lambda_n}\|}$.*

(ii) there exists $A, B > 0$ such that

$$A\|L^*f\|^2 \leq \sum_{n=1}^{\infty}\frac{|f(\lambda_n)|^2}{\|K_{\lambda_n}\|^2} \leq B\|f\|^2, \quad \text{for any } f \in \mathcal{H};$$

(iii) $\left\{\dfrac{K_{\lambda_n}}{\|K_{\lambda_n}\|}\right\}_{n=1}^{\infty}$ *is a Bessel sequence and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$Lf = \sum_{n=1}^{\infty} \langle f, g_n \rangle \frac{K_{\lambda_n}}{\|K_{\lambda_n}\|};$$

(iv) $\displaystyle\sum_{n=1}^{\infty} \frac{|f(\lambda_n)|^2}{\|K_{\lambda_n}\|^2} < \infty$ *and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$L^* f = \sum_{n=1}^{\infty} \frac{f(\lambda_n)}{\|K_{\lambda_n}\|} g_n, \forall f \in \mathcal{H}.$$

DEFINITION 4 *We say that a sequence of distinct points* $\{\lambda_n\}_{n=1}^{\infty}$ *in* Ω *is a sampling sequence for* L *if there exists two positive constants* A, B *such that*

$$A\|L^* f\|_{\mathcal{H}}^2 \leq \sum_{n=1}^{\infty} \frac{|f(\lambda_n)|^2}{\|K_{\lambda_n}\|^2} \leq B\|f\|_{\mathcal{H}}^2 \tag{2.1}$$

In the case when L is the identity operator, we have the notion of sampling sequence for $\mathcal{H}$. The normalized kernel is given by the following relation:

$$k_\lambda(z) := \frac{K_\lambda(z)}{\|K_\lambda\|}.$$

REMARK 2 *The relation (ii) in Theorem 4 is equivalent with the fact that* $\{k_{\lambda_n}\}_{n=1}^{\infty}$ *is a frame for* L*:*

$$A\|L^* f\|_{\mathcal{H}}^2 \leq \sum_{n=1}^{\infty} |\langle f, k_{\lambda_n} \rangle|^2 \leq B\|f\|_{\mathcal{H}}^2$$

because

$$\begin{aligned} \langle f, k_{\lambda_n} \rangle &= \langle f, \frac{K_{\lambda_n}}{\|K_{\lambda_n}\|_{\mathcal{H}}} \rangle \\ &= \frac{1}{\|K_{\lambda_n}\|_{\mathcal{H}}} \langle f, K_{\lambda_n} \rangle \\ &= \frac{1}{\|K_{\lambda_n}\|_{\mathcal{H}}} f(\lambda_n). \end{aligned}$$

3 ATOMIC DECOMPOSITIONS FOR OPERATORS IN BERGMAN SPACES

Let $\mathbb{C}$ be the complex plane and let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the open unit disk in $\mathbb{C}$. We denote by $dA(z)$ the area measure on $\mathbb{D}$, normalized such that the area measure on $\mathbb{D}$ is 1 :

$$dA(z) = \frac{1}{\pi} dxdy = \frac{1}{\pi} rdrd\theta, \quad z = x + iy = re^{i\theta}.$$

The Bergman space L^2_a (sometimes denoted by A^2) is the subset of $L^2(\mathbb{D})$ consisting of analytic functions. L^2_a is a reproducing kernel Hilbert space, with reproducing kernel :

$$K(z, \lambda) = K_\lambda(z) = \frac{1}{(1 - \overline{\lambda}z)^2}.$$

and normalized kernel

$$k_\lambda(z) = \frac{K_\lambda(z)}{\|K_\lambda\|} = \frac{1 - |\lambda|^2}{(1 - \overline{\lambda}z)^2}, \quad \lambda \in \mathbb{D}.$$

For a detailed account results of atomic decompositions for the identity operator on Bergman spaces see [19].

Then, from Theorem 4, we obtain the following Corollary:

COROLLARY 5 *Let $\{\lambda_n\}_{n=1}^\infty \subset \mathbb{D}$. The following are equivalent*

(i) $\{k_{\lambda_n}\}_{n=1}^\infty$ is an atomic system for L i.e.

- *the series $\sum_{n=1}^\infty c_n \frac{1 - |\lambda_n|^2}{(1 - \overline{\lambda}_n z)^2}$ converges in L^2_a for all $\{c_n\} \in l^2$; and*
- *there exists a positive constant C such that for every $f \in \mathcal{H}$ there exists $a_f = (a_n) \in l^2$ such that $\|a_f\|_{l^2} \leq C\|f\|$ and $Lf(z) = \sum_{n=1}^\infty a_n \frac{1 - |\lambda_n|^2}{(1 - \overline{\lambda}_n z)^2}$;*

(ii) there exists $A, B > 0$ such that $\{\lambda_n\}_{n=1}^\infty$ is a sampling sequence for L; i.e. there exists two positive constants A, B such that

$$A\|L^* f\|^2_{\mathcal{H}} \leq \sum_{n=1}^\infty |f(\lambda_n)|^2 (1 - |\lambda_n|^2)^2 \leq B\|f\|^2_{\mathcal{H}};$$

(iii) $\{k_{\lambda_n}\}_{n=1}^\infty$ is a Bessel sequence and there exists a Bessel sequence $\{g_n\}_{n=1}^\infty$ such that

$$Lf = \sum_{n=1}^\infty \langle f, g_n \rangle \frac{1 - |\lambda_n|^2}{(1 - \overline{\lambda}_n z)^2};$$

(iv) $\sum_{n=1}^{\infty}(1-|\lambda_n|^2)^2|f(\lambda_n)|^2 < \infty$ *and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$L^*f = \sum_{n=1}^{\infty}(1-|\lambda_n|^2)f(\lambda_n)g_n.$$

More general, for $\eta > -1$ let dA_η be the area measure on $\mathbb{D}$,

$$dA_\eta(z) = (\eta+1)(1-|z|^2)^\eta dA(z).$$

The standard weighted Bergman space $L^2_a(dA_\eta)$ is the subset of $L^2(\mathbb{D}, dA_\eta)$, consisting of analytic functions.

$$L^2(\mathbb{D}, dA_\eta) = \{f : \mathbb{D} \to \mathbb{C} : \|f\|^2_{2,\eta} := \int_{\mathbb{D}} |f(z)|^2 dA_\eta < \infty\}.$$

Then $L^2_a(dA_\eta)$ is a RKHS with reproducing kernel

$$K^\eta(z,\lambda) = K^\eta_\lambda(z) = \frac{1}{(1-\overline{\lambda}z)^{2+\eta}}, \quad z, \lambda \in \mathbb{D}$$

and normalized reproducing kernel

$$k^\eta_\lambda(z) = \frac{K^\eta_\lambda(z)}{\|K^\eta_\lambda\|_{2,\eta}} = \frac{(1-|\lambda|^2)^{1+\frac{\eta}{2}}}{(1-\overline{\lambda}z)^{2+\eta}}, \quad z, \lambda \in \mathbb{D}.$$

COROLLARY 6 *Let* $\{\lambda_n\}_{n=1}^{\infty} \subset \mathbb{D}$. *The following are equivalent*

(i) $\{k^\eta_{\lambda_n}\}_{n=1}^{\infty}$ *is an atomic system for* L *i.e.*

- *the series* $\sum_{n=1}^{\infty} c_n \frac{(1-|\lambda_n|^2)^{1+\frac{\eta}{2}}}{(1-\overline{\lambda}_n z)^{2+\eta}}$ *converges in* $L^2_a(dA_\eta)$ *for all* $\{c_n\} \in l^2$; *and*
- *there exists* $C > 0$ *such that for every* $f \in \mathcal{H}$ *there exists* $a_f = (a_n) \in l^2$ *such that* $\|a_f\|_{l^2} \le C\|f\|$ *and* $Lf(z) = \sum_{n=1}^{\infty} a_n \frac{(1-|\lambda_n|^2)^{1+\frac{\eta}{2}}}{(1-\overline{\lambda}_n z)^{2+\eta}}$;

(ii) there exists $A, B > 0$ *such that* $\{\lambda_n\}_{n=1}^{\infty}$ *is a sampling sequence for* L; *i.e. there exists two positive constants* A, B *such that*

$$A\|L^*f\|^2 \le \sum_{n=1}^{\infty} |f(\lambda_n)|^2(1-|\lambda_n|^2)^{2+\eta} \le B\|f\|^2$$

(iii) $\{k^\eta_{\lambda_n}\}_{n=1}^{\infty}$ *is a Bessel sequence and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$Lf = \sum_{n=1}^{\infty} \langle f, g_n\rangle \frac{(1-|\lambda_n|^2)^{1+\frac{\eta}{2}}}{(1-\overline{\lambda}_n z)^{2+\eta}};$$

(iv) $\sum_{n=1}^{\infty}(1-|\lambda_n|^2)^{2+\eta}|f(\lambda_n)|^2 < \infty$ *and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$L^*f = \sum_{n=1}^{\infty}(1-|\lambda_n|^2)^{1+\frac{\eta}{2}} f(\lambda_n) g_n.$$

REMARK 3 *The above Corollary is a generalization of some results in [18].*

The weighted Bergman space $A^2(\omega) = L^2_a(\omega)$ is the subset $L^2(\omega)$, consisting of analytic functions.

$$L^2(\omega) = \{f : \mathbb{D} \to \mathbb{C} : \|f\|^2_{A^2(\omega)} := \int_{\mathbb{D}} |f(z)|^2 \omega(z) dA(z) < \infty\}.$$

For $\eta > -1$ the class $B_2(\eta)$ consists of weights ω with the property that there exists a constant $c > 0$ such that

$$\left(\int_{S(\theta,h)} \omega dA_\eta\right)\left(\int_{S(\theta,h)} \omega^{-1} dA_\eta\right) \leq c[A_\eta(S(\theta,h))]^2$$

for any Carleson square:

$$S(\theta,h) = \left\{z = re^{i\alpha} : 1-h < r < 1, |\theta-\alpha| < \frac{h}{2}\right\}, \quad \theta \in [0,2\pi], h \in (0,1).$$

The weights ω considered here are called Békollé weights.

In the following, we consider $\alpha \in (0,1)$, $\eta > -1$ and $\dfrac{\omega}{(1-|z|^2)^\eta} \in B_2(\eta)$. Then $L^2_a(\omega)$ is a reproducing kernel Hilbert space. We denote by

$$K^{\eta,\omega}(z,\lambda) = K^{\eta,\omega}_\lambda(z)$$

the reproducing kernel of this space.

In [6], O. Constantin gave the following estimation for the norm of Bergman kernel $K^{\eta,\omega}_\lambda$:

$$\|K^{\eta,\omega}_\lambda\|^2 \sim \left(\int_{D_{\lambda,\alpha}} \omega dA\right)^{-1},$$

for every disc $D_{\lambda,\alpha} = \{z \in \mathbb{D} : |z-\lambda| < \alpha(1-|\lambda|)\}$.
By $\sim$ we mean that the involved constants are independent of $\lambda \in \mathbb{D}$. (For two real valued functions E_1, E_2 we write $E_1 \sim E_2$ if there there exists a positive constant c independent of the argument such that $\dfrac{1}{c}E_1 \leq E_2 \leq cE_1$).

COROLLARY 7 *Let* $\{\lambda_n\}_{n=1}^{\infty} \subset \mathbb{D}$. *Then the following statements are equivalent*

(i) $\left\{K_{\lambda_n}^{\eta,\omega}\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}\right\}_{n=1}^{\infty}$ *is an atomic system for* L*, i.e.*

- *the series* $\sum_{n=1}^{\infty} c_n\left\{K_{\lambda_n}^{\eta,\omega}\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}\right\}$ *converges in* $L_a^2(\omega)$ *for all* $\{c_n\}\in l^2$*; and*
- *there exists* $C>0$ *such that for every* $f\in\mathcal{H}$ *there exists* $a_f=(a_n)\in l^2$ *such that* $\|a_f\|_{l^2}\le C\|f\|$ *and* $Lf(z)=\sum_{n=1}^{\infty} a_n\left\{K_{\lambda_n}^{\eta,\omega}\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}\right\}$;

(ii) there exists $A,B>0$ *such that* $\{\lambda_n\}_{n=1}^{\infty}$ *is a sampling sequence for* L*; i.e. there exists two positive constants* A,B *such that*

$$A\|L^*f\|_{\mathcal{H}}^2\le\sum_{n=1}^{\infty}|f(\lambda_n)|^2\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)\le B\|f\|_{\mathcal{H}}^2;$$

(iii) $\left\{K_{\lambda_n}^{\eta,\omega}\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}\right\}_{n=1}^{\infty}$ *is a Bessel sequence and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$Lf=\sum_{n=1}^{\infty}\langle f,g_n\rangle\left\{K_{\lambda_n}^{\eta,\omega}\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}\right\};$$

(iv) $\sum_{n=1}^{\infty}|f(\lambda_n)|^2\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)<\infty$ *and there exists a Bessel sequence* $\{g_n\}_{n=1}^{\infty}$ *such that*

$$L^*f=\sum_{n=1}^{\infty}f(\lambda_n)\left(\int_{D_{\lambda_n,\alpha}}\omega dA\right)^{1/2}g_n.$$

The above Corollary provides a generalization of Theorem 2.3 in [2]. Also, our proof is more simple, with a different technique.

4 ATOMIC DECOMPOSITIONS FOR OPERATORS IN FOCK SPACES

Let $dA(z)$ be the usual Lebesgue measure on $\mathbb{C}$. For $\alpha>0$, $dA_\alpha(z)$ is defined as follows

$$dA_\alpha(z)=\frac{\alpha}{\pi}e^{-\alpha|z|^2}dA(z).$$

The Fock space F_α^2 is the space of all entire functions f on $\mathbb{C}$ for which

$$\|f\|_{F_\alpha^2}^2=\int_{\mathbb{C}}|f(z)|^2dA_\alpha(z)<\infty.$$

F^2_α is a RKHS, with reproducing kernel

$$K(z,\lambda) = K_\lambda(z) = e^{\alpha z\overline{\lambda}}$$

and normalized reproducing kernel

$$k^\alpha_\lambda(z) = \frac{K_\lambda(z)}{\|K_\lambda\|} = \frac{e^{\alpha z|\overline{\lambda}}}{e^{\frac{\alpha|\lambda|^2}{2}}} = e^{\alpha(z\overline{\lambda}-\frac{|\lambda|^2}{2})}.$$

For details on Fock spaces see the book of K. Zhu [20]. See also [17]. Atomic decompositions for the identity operator on Fock spaces were obtained in [14].

COROLLARY 8 *Let $\{\lambda_n\}_{n=1}^\infty \subset \mathbb{D}$. The following are equivalent*

(i) $\{k^\alpha_{\lambda_n}\}_{n=1}^\infty$ is an atomic system for L i.e.

- *the series $\sum_{n=1}^\infty c_n e^{\alpha(z\overline{\lambda}_n-\frac{|\lambda_n|^2}{2})}$ converges in F^2_α for all $\{c_n\} \in l^2$; and*
- *there exists $C > 0$ such that for every $f \in \mathcal{H}$ there exists $a_f = (a_n) \in l^2$ such that $\|a_f\|_{l^2} \leq C\|f\|$ and $Lf(z) = \sum_{n=1}^\infty a_n e^{\alpha(z\overline{\lambda}_n-\frac{|\lambda_n|^2}{2})}$.*

(ii) there exists $A, B > 0$ such that $\{\lambda_n\}_{n=1}^\infty$ is a sampling sequence for L; i.e. there exists two positive constants A, B such that

$$A\|L^*f\|^2 \leq \sum_{n=1}^\infty |f(\lambda_n)|^2 e^{-\alpha|\lambda_n|^2} \leq B\|f\|^2;$$

(iii) $\{k^\alpha_{\lambda_n}\}_{n=1}^\infty$ is a Bessel sequence and there exists a Bessel sequence $\{g_n\}_{n=1}^\infty$ such that

$$Lf = \sum_{n=1}^\infty \langle f, g_n\rangle e^{\alpha(z\overline{\lambda}_n-\frac{|\lambda_n|^2}{2})};$$

(iv) $\sum_{n=1}^\infty e^{-\alpha|\lambda_n|^2}|f(\lambda_n)|^2 < \infty$ and there exists a Bessel sequence $\{g_n\}_{n=1}^\infty$ such that

$$L^*f = \sum_{n=1}^\infty e^{-\alpha\frac{|\lambda_n|^2}{2}} f(\lambda_n) g_n.$$

REMARK 4 *The above Corollary is a generalization of some results in [16].*

Acknowledgements. I would like to thanks Dr. Olivia Constantin for introducing me in the theory of Bergman and Fock spaces and also for the careful reading of this paper and her useful comments.

REFERENCES

[1] P.G. Casazza, The art of frame theory, Taiwanese J.Math., 4(2)(2000) 129-202.

[2] G. Chacón, Toeplitz Operators on Weighted Bergman Spaces, arXiv.org:1208.2917, 1–9.

[3] O. Christensen, An Introduction to Frames and Riesz Bases, Birkhaüser, 2003.

[4] C.K. Chui. An Introduction to Wavelets. Academic Press, San Diego, CA, 1992.

[5] R.R. Coifman, R. Rochberg, Representation theorems for holomorphic functions in L^p, Asterisque 77 (1980), 11–66 .

[6] O. Constantin, Carleson embeddings and some classes of operators on weighted Bergman spaces, J. Math. Anal. Appl. 365 (2010) 668–682.

[7] L. Găvruţa, Frames for operators, Appl. Comput. Harmon. Anal. 32 (2012) 139–144.

[8] I. Daubechies, Ten Lectures on Wavelets. Philadelphia, PA: SIAM, 1992.

[9] I. Daubechies, A. Grossmann and Y. Meyer, Painless nonorthogonal expansions. J.Math.Phys. 27(1986) 1271-1283.

[10] R.J. Duffin, A.C. Schaeffer, A class of nonharmonic Fourier series, Trans.Amer.Math.Soc.72(1952) 341–366.

[11] H.G. Feichtinger and T. Werther, Atomic systems for subspaces, Proceedings SampTA 2001 (L.Zayed, ed.), Orlando, FL, 2001, pp. 163-165.

[12] K. Gröchenig, Foundations of Time-Frequency Analysis, Birkhäuser, Boston, 2001.

[13] C. Heil, A Basis Theory Primer, Birkhäuser, Boston, 2010.

[14] S. Janson, J. Peetre, R. Rochberg, Hankel forms and the Fock space, Revista Mat. Iberoamericana, 3 (1987), 61–129.

[15] J. Kovačević and A. Chebira, An Introduction to Frames, Foundations and Trends in Signal Processing,2, No. 1 (2008) 1-94.

[16] M. Tatari, S.M. Vaezpour, A. Pishinian, On Some Properties of Fock Space F^2_α by Frame Theory, Int. J. Contemp. Sciences, Vol. 5, 2010, no. 23, 1107–1114.

[17] Y.J. Tung, Fock spaces, Ph.D. Thesis, University of Michigan, 2005.

[18] S.M. Vaezpour, M. Tatari, A. Pishinian, Atomic Decomposition for Weighted Bergman Space L_a^2 on the Unit Ball of C, Int. J. Contemp. Math. Sciences, Vol. 4, 2009, no. 17, 827–831.

[19] K. Zhu, Operator Theory in Function Spaces, second ed., Math. Surveys Monogr., vol. 138, Amer. Math. Soc., Providence, RI, 2007.

[20] K. Zhu, Analysis on Fock Spaces, Springer-Verlag, New York, 2012.

Politehnica University of Timişoara,
Department of Mathematics
Piaţa Victoriei no. 2, 300006 Timişoara, Romania

Fakultät fur Mathematik, University of Viena
Oskar-Morgenstern-Platz 1 1090 Wien, Austria
E-mail: gavruta_laura@yahoo.com